

The $B(B_s) \rightarrow D_{(s)}P$, $D_{(s)}V$, $D_{(s)}^*P$ and $D_{(s)}^*V$ decays in the perturbative QCD approach

Run-Hui Li^{1,2}, Cai-Dian Lü¹, and Hao Zou¹

¹*Institute of High Energy Physics, P.O. Box 918(4), Beijing 100049, China and*

²*Department of Physics, Shandong University, Jinan 250100, China*

(Dated: March 27, 2008)

Two-body non-leptonic charmed decays $B_{(s)} \rightarrow D_{(s)}P$, $D_{(s)}^*P$, $D_{(s)}V$ and $D_{(s)}^*V$ are analyzed in perturbative QCD approach, where P and V denote the light pseudoscalar meson and vector meson, respectively. We test the D meson wave function by a χ^2 fit with experimental data of six $B \rightarrow DP$ channels. We give the branching ratios of all the charmed B decay channels, most of which agree with experiments amazingly well. The predicted B_s decays can be confronted with the future experimental data. By straightforward calculations, our pQCD approach gives the right relative strong phase of a_2/a_1 with experiments. We also predict the percentage of transverse polarizations in $B_{(s)} \rightarrow D^*V$ decay channels.

I. INTRODUCTION

B physics experiments provide a good test of the standard model and severe constraints of new physics parameters. Recent years more and more efforts have been made to the study of B meson decays both experimentally and theoretically. In the near future, there will be more and more data in B physics thanks to the run of B factories, Tevatron and LHCb. Theoretically, a great improvement has been made to the study of exclusive decays of B mesons. In the history, naive factorization [1] is a successful method to explain many decay branching ratios [2], but it failed to explain color-suppressed processes such as $\bar{B}^0 \rightarrow D^0\pi^0$ [3]. Currently, perturbative QCD factorization approach (pQCD) [4] is one of the popular methods to deal with the two-body non-leptonic decays of B mesons. It explains the experiments successfully, especially for the direct CP asymmetries [5] when the final states are two light mesons, which inspires people to see how far it will go.

Charmed decays of B (B_s) mesons are more complicated than the decays with only light mesons as final states. The $B \rightarrow D$ transitions involve three scales: M_B , M_D , $\bar{\Lambda}$, respectively. The factorization was proved in soft-collinear effective theory [6] with less predictions than pQCD approach, since they need more inputs than pQCD approach. In $B \rightarrow$ light transition, the light spectator quark in B meson is soft, while it is collinear in the final state meson, so that a hard gluon




FIG. 1: color allowed diagrams in pQCD approach for $B \rightarrow DP$ decays

is needed to connect it to the four quark operator shown in Figure 1. The momentum square of the hard gluon connecting the spectator quark in $B \rightarrow D$ transition is only a factor of $(1 - m_D^2/m_B^2)$ than the $B \rightarrow \text{light}$ transition, which shows pQCD should also work well in $B \rightarrow D$ transitions. The hierarchy: $M_B \gg M_D \gg \bar{\Lambda}$ is used in pQCD framework [7]. Some separate calculations on $B \rightarrow D$ decays in pQCD approach are carried on [8, 9] in the leading order of M_D/M_B and Λ/M_B expansions. It's found that the pQCD do work well since the D meson recoils fast.

In this paper, we calculate all the processes of a $B_{(s)}$ meson decays to a $D_{(s)}^{(*)}$ meson and a light pseudoscalar meson or vector meson. Only tree diagrams contribute to these processes involving only one kind of CKM matrix elements. So there's no direct CP asymmetry in these decays. The light cone distribution amplitudes (LCDA) of mesons are necessary inputs in the pQCD framework. The light mesons' distribution amplitudes (DAs) have been well studied and calculated by QCD sum rules. But there are little studies on heavy mesons' DAs, especially for D meson. In this paper we collect several candidate distribution amplitude models for D meson, then we fit out the parameters using the experimental results and make a comparison among them.

The paper is organized as follows: Section II contains the conventions and notations that we adopt, together with all the wave functions used in this paper. The pQCD analytic formulae for the amplitudes are given in section III. Section IV contain the numerical results and discussions. Section V is our summary.

II. ANALYTIC FORMULAE

For the charmed B decays we considered, only the tree operators of the standard effective weak Hamiltonian contribute. The Hamiltonian is given by:

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} V_{cb} V_{uq'}^* [C_1(\mu) O_1(\mu) + C_2(\mu) O_2(\mu)] \quad . \quad (1)$$

The tree operators are

$$O_1 = (\bar{c}_\alpha b_\beta)_{V-A} (\bar{q}'_\beta u_\alpha)_{V-A} , O_2 = (\bar{c}_\alpha b_\alpha)_{V-A} (\bar{q}'_\beta u_\beta)_{V-A} , \quad (2)$$

with $q' = d, s$. $\alpha \beta$ are the color indexes, and $(\bar{q}_1 q_2)_{V-A} \equiv \bar{q}_1 \gamma^\mu (1 - \gamma^5) q_2$. The $V_{q_1 q_2}$ in the Hamiltonian denote the CKM matrix elements.

In the factorization formulae, the combinations of Wilson coefficients usually appear. In this paper, they are defined as follows:

$$a_1 = C_2 + C_1/3 , a_2 = C_1 + C_2/3 . \quad (3)$$

In this paper, all the momentum are defined with the light cone coordinate. Two light-like vectors n and v are defined, with $n^2 = 0$, $v^2 = 0$ and $n \cdot v = 1$. n can be written as $(1, 0, \mathbf{0}_T)$, and v is $(0, 1, \mathbf{0}_T)$. The momentum of B meson, D meson and the lightest meson are denoted as P_1 , P_2 and P_3 respectively. At the rest frame of B meson, the light meson moves very fast. So P_3^+ or P_3^- can be treated as zero. In this paper, the momentum are defined as:

$$P_1 = \frac{M_B}{\sqrt{2}}(1, 1, \mathbf{0}_T) , P_2 = \frac{M_B}{\sqrt{2}}(1, r^2, \mathbf{0}_T) , P_3 = \frac{M_B}{\sqrt{2}}(0, 1 - r^2, \mathbf{0}_T) . \quad (4)$$

The momentum of the light anti-quark in the B meson and D meson are denoted as k_1 and k_2 respectively, while k_3 is defined as the quark momentum of lightest meson. In our calculation they are taken as:

$$\begin{aligned} k_2 &= (x_2 \frac{M_B}{\sqrt{2}}, 0, \mathbf{k}_{2\perp}) , k_3 = (0, x_3 \frac{(1 - r^2)M_B}{\sqrt{2}}, \mathbf{k}_{3\perp}) , \\ k_1 &= (x_1 \frac{M_B}{\sqrt{2}}, 0, \mathbf{k}_{1\perp}) \text{ for color suppressed contributions,} \\ k_1 &= (0, x_1 \frac{M_B}{\sqrt{2}}, \mathbf{k}_{1\perp}) \text{ for the others.} \end{aligned} \quad (5)$$

with x_1 , x_2 and x_3 as the momentum fraction, and $\mathbf{k}_{1\perp}$, $\mathbf{k}_{2\perp}$ and $\mathbf{k}_{3\perp}$ as the transverse momentum of the quark.

A. Wave functions of $B_{(s)}$ mesons

In pQCD calculation, the light-cone wave functions of the mesons are needed. The B meson and B_s meson have the similar structure of wave function, except different values of parameters characterizing a small SU(3) breaking effect. In general, the $B_{(s)}$ meson wave function are always

decomposed into the following Lorentz structures:

$$\begin{aligned} & \int \frac{d^4 z}{(2\pi)^4} e^{ik_1 \cdot z} \langle 0 | \bar{b}_\alpha(0) d_\beta(z) | B_{(s)}(P_1) \rangle \\ &= \frac{i}{\sqrt{2N_c}} \left\{ (\not{P}_1 + M_{B_{(s)}}) \gamma_5 [\phi_{B_{(s)}}(k_1) - \frac{\not{h} - \not{b}}{\sqrt{2}} \bar{\phi}_{B_{(s)}}(k_1)] \right\}_{\beta\alpha}. \end{aligned} \quad (6)$$

There are two distribution amplitudes in the above equation. However, $\bar{\phi}_{B_{(s)}}(k_1)$ gives smaller contribution [10]. We will neglect it in our calculation and only keep the first Lorentz structure.

$$\Phi_{B_{(s)}} = \frac{i}{\sqrt{2N_c}} (\not{P}_1 + M_{B_{(s)}}) \gamma_5 \phi_{B_{(s)}}(k_1). \quad (7)$$

The distribution amplitude in the b-space is:

$$\phi_{B_{(s)}}(x, b) = N_{B_{(s)}} x^2 (1-x)^2 \exp \left[-\frac{1}{2} \left(\frac{x M_{B_{(s)}}}{\omega_b} \right)^2 - \frac{\omega_b^2 b^2}{2} \right], \quad (8)$$

with b as the conjugate space coordinate of $\mathbf{k}_{1\perp}$. $N_{B_{(s)}}$ is the normalization constant, which is determined by the normalization condition:

$$\int_0^1 dx \phi_{B_{(s)}}(x, b=0) = \frac{f_{B_{(s)}}}{2\sqrt{2N_c}}. \quad (9)$$

For parameter ω_b , we take the value 0.40 GeV for B_d^0 and $B^\pm$ mesons, and 0.50 ± 0.05 GeV for B_s^0 meson, characterizing the small SU(3) breaking effect as argued in [11].

B. Wave functions of light pseudoscalar mesons

The decay constant of the pseudoscalar mesons is defined by:

$$\langle 0 | \bar{q}_1 \gamma_\mu \gamma_5 q_2 | P(P_3) \rangle = i f_P P_\mu. \quad (10)$$

The decay constant for π and K are $f_\pi = 131$ MeV, $f_K = 160$ MeV.

The light cone distribution amplitudes (for out-going state) for light pseudoscalar mesons is:

$$\begin{aligned} & \langle P(P_3) | \bar{q}_{2\beta}(z) q_{1\alpha}(0) | 0 \rangle \\ &= -\frac{i}{\sqrt{2N_c}} \int_0^1 dx e^{ixP \cdot z} \left[\gamma_5 \not{P}_3 \phi^A(x) + \gamma_5 m_0 \phi^P(x) + m_0 \gamma_5 (\not{b} \not{h} - 1) \phi^T(x) \right]_{\alpha\beta}, \end{aligned} \quad (11)$$

where v is the light cone direction that the light pseudoscalar meson's momentum is defined on.

The chiral scale parameter m_0 is defined as $m_0 = \frac{M_P^2}{m_{q1} + m_{q2}}$.

The distribution amplitudes are usually expanded by the Gegenbauer polynomials. Their expressions are:

$$\phi_P^A(x) = \frac{3f_P}{\sqrt{2N_c}}x(1-x) \left[1 + a_1^A C_1^{3/2}(t) + a_2^A C_2^{3/2}(t) + a_4^A C_4^{3/2}(t) \right], \quad (12)$$

$$\phi_P^p(x) = \frac{f_P}{2\sqrt{2N_c}} \left[1 + a_2^p C_2^{1/2}(t) + a_4^p C_4^{1/2}(t) \right], \quad (13)$$

$$\phi_P^T(x) = -\frac{f_P}{2\sqrt{2N_c}} \left[C_1^{1/2}(t) + a_3^T C_3^{1/2}(t) \right], \quad (14)$$

with $t=2x-1$. The coefficients of the Gegenbauer polynomials are [26]

$$\begin{aligned} a_{2\pi}^A &= 0.44, \quad a_{4\pi}^A = 0.25, \quad a_{1K}^A = 0.17, \quad a_{2K}^A = 0.2, \\ a_{2\pi}^p &= 0.43, \quad a_{4\pi}^p = 0.09, \quad a_{2K}^p = 0.24, \quad a_{4K}^p = -0.11, \\ a_{3\pi}^T &= 0.55, \quad a_{3K}^T = 0.35. \end{aligned} \quad (15)$$

For η and η' , the mixing mechanism must be taken into consideration. We will take the method presented in ref. [12], where the η_n and η_s are chosen as the basis of mixing:

$$\begin{pmatrix} |\eta\rangle \\ |\eta'\rangle \end{pmatrix} = U(\phi) \begin{pmatrix} |\eta_n\rangle \\ |\eta_s\rangle \end{pmatrix}, \quad (16)$$

with

$$|\eta_n\rangle = \frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d), \quad |\eta_s\rangle = \bar{s}s, \quad (17)$$

$$U(\phi) = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}, \quad (18)$$

where the mixing angle $\phi = 39.3^\circ \pm 1.0^\circ$.

The assumption that the distribution amplitudes of η_n and η_s is the same as the distribution amplitudes of π is adopted, except different decay constants and chiral parameters. The decay constant of η_n and η_s is taken from ref.[12]:

$$f_n = (1.07 \pm 0.02)f_\pi = (139.1 \pm 2.6) \text{ MeV}, \quad f_s = (1.34 \pm 0.06)f_\pi = (174.2 \pm 7.8) \text{ MeV}. \quad (19)$$

And the chiral parameters are given by

$$m_0^{\bar{n}n} = \frac{1}{2m_n} [m_\eta^2 \cos^2 \phi + m_{\eta'}^2 \sin^2 \phi - \frac{\sqrt{2}f_s}{f_n} (m_{\eta'}^2 - m_\eta^2) \cos \phi \sin \phi], \quad (20)$$

$$m_0^{\bar{s}s} = \frac{1}{2m_s} [m_{\eta'}^2 \cos^2 \phi + m_\eta^2 \sin^2 \phi - \frac{f_n}{\sqrt{2}f_s} (m_{\eta'}^2 - m_\eta^2) \cos \phi \sin \phi]. \quad (21)$$

TABLE I: The decay constants of vector mesons (in MeV)

f_ρ	f_{K^*}	f_ω	f_ϕ	f_ρ^T	$f_{K^*}^T$	f_ω^T	f_ϕ^T
209 ± 2	217 ± 5	195 ± 3	231 ± 4	165 ± 9	185 ± 10	151 ± 9	186 ± 9

C. Wave functions of light vector mesons

The decay constants for the vector mesons are defined by:

$$\langle 0 | \bar{q}_1 \gamma_\mu q_2 | V(p, \epsilon) \rangle = f_V m_V \epsilon_\mu, \quad \langle 0 | \bar{q}_1 \sigma_{\mu\nu} q_2 | V(p, \epsilon) \rangle = i f_V^T (\epsilon_\mu p_\nu - \epsilon_\nu p_\mu). \quad (22)$$

The longitudinal decay constant of vector mesons can be extracted experimentally [13]. And the transverse ones can be calculated by the QCD sum rule [14]. We list all of them in Table I.

The distribution amplitudes up to twist-3 of vector mesons are

$$\begin{aligned} \langle V(P, \epsilon_L^*) | \bar{q}_{2\beta}(z) q_{1\alpha}(0) | 0 \rangle &= \frac{1}{\sqrt{2N_C}} \int_0^1 dx e^{ixP \cdot z} [M_V \not{\epsilon}_L^* \phi_V(x) + \not{\epsilon}_L^* \not{P} \phi_V^t(x) + M_V \phi_V^s(x)]_{\alpha\beta}, \\ \langle V(P, \epsilon_T^*) | \bar{q}_{2\beta}(z) q_{1\alpha}(0) | 0 \rangle &= \frac{1}{\sqrt{2N_C}} \int_0^1 dx e^{ixP \cdot z} [M_V \not{\epsilon}_T^* \phi_V^v(x) + \not{\epsilon}_T^* \not{P} \phi_V^T(x) \\ &\quad + M_V i \epsilon_{\mu\nu\rho\sigma} \gamma_5 \gamma^\mu \epsilon_T^{*\nu} n^\rho v^\sigma \phi_V^a(x)]_{\alpha\beta}, \end{aligned} \quad (23)$$

where x is the momentum fraction of the q_2 quark. And convention $\epsilon^{0123} = 1$ is adopted for the Levi-Civita tensor. The twist-2 distribution amplitudes of vector mesons are expanded as:

$$\begin{aligned} \phi_V(x) &= \frac{3f_V}{\sqrt{2N_C}} x(1-x) \left[1 + a_1^\parallel C_1^{\frac{3}{2}}(t) + a_2^\parallel C_2^{\frac{3}{2}}(t) \right], \\ \phi_V^T(x) &= \frac{3f_V}{\sqrt{2N_C}} x(1-x) \left[1 + a_1^\perp C_1^{\frac{3}{2}}(t) + a_2^\perp C_2^{\frac{3}{2}}(t) \right]. \end{aligned} \quad (24)$$

We take the following values for the Gegenbauer moments [25]:

$$\begin{aligned} a_{2\rho}^\parallel &= a_{2\omega}^\parallel = 0.15 \pm 0.07, \quad a_{1K^*}^\parallel = 0.03 \pm 0.02, \quad a_{2K^*}^\parallel = 0.11 \pm 0.09, \quad a_{2\phi}^\parallel = 0.18 \pm 0.08, \\ a_{2\rho}^\perp &= a_{2\omega}^\perp = 0.14 \pm 0.06, \quad a_{1K^*}^\perp = 0.04 \pm 0.03, \quad a_{2K^*}^\perp = 0.10 \pm 0.08, \quad a_{2\phi}^\perp = 0.14 \pm 0.07. \end{aligned} \quad (25)$$

For the other distribution amplitudes, we use the asymptotic form:

$$\begin{aligned} \phi_V^t(x) &= \frac{3f_V^T}{2\sqrt{6}} t^2, \quad \phi_V^s(x) = \frac{3f_V^T}{2\sqrt{6}} (-t), \\ \phi_V^v(x) &= \frac{3f_V}{3\sqrt{6}} (1+t^2), \quad \phi_V^a(x) = \frac{3f_V}{4\sqrt{6}} (-t). \end{aligned} \quad (26)$$

D. Wave function of $D^{(*)}$ meson

The two-particle light-cone distribution amplitudes of $D^{(*)}$ meson, up to twist-3 accuracy, are defined by [7]:

$$\begin{aligned} \int \frac{d^4w}{(2\pi)^4} e^{ik \cdot w} \langle 0 | \bar{c}_\beta(0) d_\gamma(w) | D^-(P) \rangle &= -\frac{i}{\sqrt{2N_C}} [(\not{P} + M_D) \gamma_5]_{\gamma\beta} \phi_D(x), \\ \int \frac{d^4w}{(2\pi)^4} e^{ik \cdot w} \langle 0 | \bar{c}_\beta(0) d_\gamma(w) | D^{*-}(P) \rangle &= -\frac{i}{\sqrt{2N_C}} [(\not{P} + M_{D^*}) \not{\epsilon}_L \phi_{D^*}^L(x) + (\not{P} + M_{D^*}) \not{\epsilon}_T \phi_{D^*}^T(x)]_{\gamma\beta} \end{aligned} \quad (27)$$

with

$$\int_0^1 dx \phi_D(x) = \frac{f_D}{2\sqrt{2N_C}}, \quad \int_0^1 dx \phi_{D^*}^L(x) = \frac{f_{D^*}}{2\sqrt{2N_C}}, \quad \int_0^1 dx \phi_{D^*}^T(x) = \frac{f_{D^*}^T}{2\sqrt{2N_C}}, \quad (28)$$

as the normalization conditions. In the heavy quark limit we have

$$f_{D^*}^T - f_{D^*} \frac{m_c + m_d}{M_{D^*}} \sim f_{D^*} - f_{D^*}^T \frac{m_c + m_d}{M_{D^*}} \sim O(\bar{\Lambda}/M_{D^*}). \quad (29)$$

Thus we use $f_{D^*}^T = f_{D^*}$ in the calculation. There are several candidate distribution amplitude models for D meson. We collect as below:

$$\begin{aligned} \phi_D^{(Gen)}(x, b) &= \frac{1}{2\sqrt{2N_C}} f_D 6x(1-x)[1 + C_D(1-2x)], \\ \phi_D^{(MGen)}(x, b) &= \frac{1}{2\sqrt{2N_C}} f_D 6x(1-x)[1 + C_D(1-2x)] \exp\left[\frac{-\omega^2 b^2}{2}\right], \\ \phi_D^{(KLS)}(x, b) &= \frac{1}{2\sqrt{2N_C}} f_D N_D \sqrt{x(1-x)} \exp\left[-\frac{1}{2}\left(\frac{xM_D}{\omega}\right)^2 - \frac{\omega^2 b^2}{2}\right], \\ \phi_D^{(GN)}(x, b) &= \frac{1}{2\sqrt{2N_C}} f_D N_D x \exp\left[-\frac{xM_D}{\omega}\right] \frac{1}{1 + b^2 \omega^2}, \\ \phi_D^{(KKQT)}(x, b) &= \frac{1}{2\sqrt{2N_C}} f_D N_D x \theta(x) \theta\left(\frac{2\Lambda_D}{M_D} - x\right) J_0\left(b \sqrt{x\left(\frac{2\Lambda_D}{M_D} - x\right)}\right), \\ \phi_D^{(Huang)}(x, b) &= \frac{1}{2\sqrt{2N_C}} f_D N_D x(1-x) \exp\left[-\Lambda_D \frac{(1-x)m_d^2 + xm_c^2}{x(1-x)}\right]. \end{aligned} \quad (30)$$

In the above models x is the momentum fraction of the light quark in D meson. The first DA model $\phi_D^{(Gen)}$ was proposed in [7], which is the Gegenbauer polynomial-like form. In order to make it $k_\perp$ dependent, an exponential term is added to get $\phi_D^{(MGen)}$. The third candidate DA model $\phi_D^{(KLS)}$ was proposed in [15], which is a Gaussian type model. The fourth one [16], which is an exponential model, and the fifth model [17], which is obtained by solving the equations of motion without three-parton contributions, were first proposed for B meson. Here we use heavy quark symmetry and modify the parameters to make them D meson DAs. The sixth DA was proposed in




FIG. 2: color suppressed diagrams in pQCD approach for $B \rightarrow DP$ decays




FIG. 3: annihilation type diagrams in pQCD approach for $B \rightarrow DP$ decays

[18], which is derived from the BHL prescription [19], with $m_d = 0.35$ GeV, $m_c = 1.3$ GeV. In the above candidate DAs, only the second model has two parameters, and only $\phi_D^{(Gen)}$ and $\phi_D^{(Huang)}$ are b independent. In the next section we will try to fit out the best D meson wave function parameters with the experimental results. As for D^* meson, we just assume that $\phi_D^L = \phi_D^T = \phi_D$ according to heavy quark symmetry.

III. CALCULATION OF DECAY AMPLITUDES IN PQCD APPROACH

A. Amplitudes for $B_{(s)} \rightarrow D_{(s)}P$ decays

There are three types of diagrams that may contribute to the $B \rightarrow D^{(*)}M$ decays: color allowed diagrams (we mark this kind of contribution with the subscript *ext*) shown in Fig. 1, color suppressed diagrams (marked with *int*) shown in Fig. 2, and the annihilation type diagrams (marked with *exc*) shown in Fig. 3. And each type of diagrams contains two categories: the one with one meson can be factorized out (denoted as ξ) and the one with no meson can be factorized out (denoted as M).

The first two diagrams in Fig. 1, 2 and 3 involve only two meson wave functions, whose results

are as following:

$$\begin{aligned} \xi_{\text{ext}} = & 8\pi C_F f_P \int_0^1 dx_1 dx_2 \int_0^{1/\Lambda} b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_D(x_2) \\ & \times \left[E_e(t_e^{(1)}) h(x_1, x_2, b_1, b_2) S_t(x_2) (1 + x_2 + r) + r E_e(t_e^{(2)}) h(x_2, x_1, b_2, b_1) S_t(x_1) \right], \quad (31) \end{aligned}$$

$$\begin{aligned} \xi_{\text{int}} = & 8\pi C_F f_D \int_0^1 dx_1 dx_3 \int_0^{1/\Lambda} b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) \\ & \times \{ [(2 - x_3) \phi_P(x_3) - r_0(1 - 2x_3)(\phi_P^p(x_3) - \phi_P^T(x_3))] \\ & \times E_i(t_i^{(1)}) h(x_1, (1 - x_3)(1 - r^2), b_1, b_3) S_t(x_3) \\ & + 2r_0 \phi_P^p(x_3) E_i(t_i^{(2)}) h(1 - x_3, x_1(1 - r^2), b_3, b_1) S_t(x_1) \}, \quad (32) \end{aligned}$$

$$\begin{aligned} \xi_{\text{exc}} = & 8\pi C_F f_B \int_0^1 dx_2 dx_3 \int_0^{1/\Lambda} b_2 db_2 b_3 db_3 \phi_D(x_2) \\ & \times \left[-x_3 \phi_P(x_3) E_a(t_a^{(1)}) h_a(x_2, x_3(1 - r^2), b_2, b_3) S_t(x_3) \right. \\ & \left. + x_2 \phi_P(x_3) E_a(t_a^{(2)}) h_a(x_3, x_2(1 - r^2), b_3, b_2) S_t(x_2) \right], \quad (33) \end{aligned}$$

with the mass ratio $r_0 \equiv m_0/m_B$. f_P , f_B and f_D are the decay constants of the light pseudoscalar meson, B meson and D meson respectively. And the factor evolving with the scale t are given by:

$$\begin{aligned} E_e(t) &= \alpha_s(t) a_1(t) \exp[-S_B(t) - S_D(t)], \\ E_i(t) &= \alpha_s(t) a_2(t) \exp[-S_B(t) - S_P(t)], \\ E_a(t) &= \alpha_s(t) a_2(t) \exp[-S_D(t) - S_P(t)]. \quad (34) \end{aligned}$$

We adopt the expression of Sudakov factor for D meson as suggested in ref. [7], which is listed in appendix together with the expressions for $S_B(t)$, $S_P(t)$.

The functions h 's in the hard part of factorization formulae, derived from the factorizable diagrams, are given by

$$\begin{aligned} h(x_1, x_2, b_1, b_2) &= K_0(\sqrt{x_1 x_2} m_B b_1) \\ &\times [\theta(b_1 - b_2) K_0(\sqrt{x_2} m_B b_1) I_0(\sqrt{x_2} m_B b_2) \\ &+ \theta(b_2 - b_1) K_0(\sqrt{x_2} m_B b_2) I_0(\sqrt{x_2} m_B b_1)], \quad (35) \end{aligned}$$

$$\begin{aligned} h_a(x_2, x_3, b_2, b_3) &= \left(i\frac{\pi}{2}\right)^2 H_0^{(1)}(\sqrt{x_2 x_3} m_B b_2) \\ &\times \left[\theta(b_2 - b_3) H_0^{(1)}(\sqrt{x_3} m_B b_2) J_0(\sqrt{x_3} m_B b_3) \right. \\ &\left. + \theta(b_3 - b_2) H_0^{(1)}(\sqrt{x_3} m_B b_3) J_0(\sqrt{x_3} m_B b_2) \right], \quad (36) \end{aligned}$$

where $H^{(1)}(z) = J_0(z) + iY_0(z)$. The hard scales t are determined by

$$\begin{aligned} t_e^{(1)} &= \max(\sqrt{x_2}m_B, 1/b_1, 1/b_2), \quad t_e^{(2)} = \max(\sqrt{x_1}m_B, 1/b_1, 1/b_2), \\ t_i^{(1)} &= \max(\sqrt{(1-x_3)(1-r^2)}m_B, 1/b_1, 1/b_3), \quad t_i^{(2)} = \max(\sqrt{x_1(1-r^2)}m_B, 1/b_1, 1/b_3), \\ t_a^{(1)} &= \max(\sqrt{x_3(1-r^2)}m_B, 1/b_2, 1/b_3), \quad t_a^{(2)} = \max(\sqrt{x_2(1-r^2)}m_B, 1/b_2, 1/b_3). \end{aligned} \quad (37)$$

The formulae for the last two diagrams in Fig. 1, Fig. 2 and Fig. 3 contain the kinematics variables of the three mesons. Their expressions are:

$$\begin{aligned} \mathcal{M}_{\text{ext}} &= 16\pi\sqrt{2N_c}C_F \int_0^1 [dx] \int_0^{1/\Lambda} b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) \phi_D(x_2) \phi_P(x_3) \\ &\quad \times \left[x_3 E_b(t_b^{(1)}) h_b^{(1)}(x_i, b_i) - (1-x_3+x_2) E_b(t_b^{(2)}) h_b^{(2)}(x_i, b_i) \right], \end{aligned} \quad (38)$$

$$\begin{aligned} \mathcal{M}_{\text{int}} &= 16\pi\sqrt{2N_c}C_F \int_0^1 [dx] \int_0^{1/\Lambda} b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_D(x_2) \\ &\quad \times \left[((x_3-1-x_2)\phi_P(x_3) + r_0(1-x_3)(\phi_P^p(x_3) - \phi_P^T(x_3))) E_d(t_d^{(1)}) h_d^{(1)}(x_i, b_i) \right. \\ &\quad \left. + [(1-x_2)\phi_P(x_3) + r_0(x_3-1)(\phi_P^p(x_3) + \phi_P^T(x_3))] E_d(t_d^{(2)}) h_d^{(2)}(x_i, b_i) \right], \end{aligned} \quad (39)$$

$$\begin{aligned} \mathcal{M}_{\text{exc}} &= 16\pi\sqrt{2N_c}C_F \int_0^1 [dx] \int_0^{1/\Lambda} b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_D(x_2) \\ &\quad \times \left[x_3 \phi_P(x_3) E_f(t_f^{(1)}) h_f^{(1)}(x_i, b_i) - x_2 \phi_P(x_3) E_f(t_f^{(2)}) h_f^{(2)}(x_i, b_i) \right], \end{aligned} \quad (40)$$

with $[dx] \equiv dx_1 dx_2 dx_3$. The expressions for the evolution factors are

$$\begin{aligned} E_b(t) &= \alpha_s(t) \frac{C_1(t)}{N_C} \exp[-S(t)|_{b_2=b_1}], \\ E_d(t) &= \alpha_s(t) \frac{C_2(t)}{N_C} \exp[-S(t)|_{b_3=b_1}], \\ E_f(t) &= \alpha_s(t) \frac{C_2(t)}{N_C} \exp[-S(t)|_{b_3=b_2}], \end{aligned} \quad (41)$$

with the Sudakov exponent $S = S_B + S_D + S_P$.

The functions $h^{(j)}$, $j = 1$ and 2 , in these amplitudes are

$$h_b^{(j)} = [\theta(b_1 - b_3)K_0(Bm_B b_1)I_0(Bm_B b_3) + \theta(b_3 - b_1)K_0(Bm_B b_3)I_0(Bm_B b_1)] \\ \times \left\{ \begin{array}{ll} K_0(B_j m_B b_3) & \text{for } B_j^2 \geq 0 \\ \frac{i\pi}{2}H_0^{(1)}(\sqrt{|B_j^2|}m_B b_3) & \text{for } B_j^2 \leq 0 \end{array} \right\}, \quad (42)$$

$$h_d^{(j)} = [\theta(b_1 - b_2)K_0(Dm_B b_1)I_0(Dm_B b_2) + \theta(b_2 - b_1)K_0(Dm_B b_2)I_0(Dm_B b_1)] \\ \times \left\{ \begin{array}{ll} K_0(D_j m_B b_2) & \text{for } D_j^2 \geq 0 \\ \frac{i\pi}{2}H_0^{(1)}(\sqrt{|D_j^2|}m_B b_2) & \text{for } D_j^2 \leq 0 \end{array} \right\}, \quad (43)$$

$$h_f^{(j)} = i\frac{\pi}{2} \left[\theta(b_1 - b_2)H_0^{(1)}(Fm_B b_1)J_0(Fm_B b_2) + \theta(b_2 - b_1)H_0^{(1)}(Fm_B b_2)J_0(Fm_B b_1) \right] \\ \times \left\{ \begin{array}{ll} K_0(F_j m_B b_1) & \text{for } F_j^2 \geq 0 \\ \frac{i\pi}{2}H_0^{(1)}(\sqrt{|F_j^2|}m_B b_1) & \text{for } F_j^2 \leq 0 \end{array} \right\}, \quad (44)$$

with the variables

$$\begin{aligned} B^2 &= x_1 x_2, \\ B_1^2 &= x_1 x_2 - x_2 x_3 (1 - r^2), \\ B_2^2 &= x_1 x_2 - x_2 (1 - x_3) (1 - r^2), \\ D^2 &= x_1 (1 - x_3) (1 - r^2), \\ D_1^2 &= (x_1 - x_2) (1 - x_3) (1 - r^2), \\ D_2^2 &= (x_1 + x_2) r^2 - (1 - x_1 - x_2) (1 - x_3) (1 - r^2), \\ F^2 &= x_2 x_3 (1 - r^2), \\ F_1^2 &= x_2 (x_1 - x_3 (1 - r^2)), \\ F_2^2 &= 1 - (1 - x_2) (1 - x_1 - x_3 (1 - r^2)). \end{aligned} \quad (45)$$

The scales $t^{(j)}$ are given by

$$\begin{aligned} t_b^{(j)} &= \max(Bm_B, \sqrt{|B_j^2|}m_B, 1/b_1, 1/b_3), \\ t_d^{(j)} &= \max(Dm_B, \sqrt{|D_j^2|}m_B, 1/b_1, 1/b_2), \\ t_f^{(j)} &= \max(Fm_B, \sqrt{|F_j^2|}m_B, 1/b_1, 1/b_2). \end{aligned} \quad (46)$$

The decay amplitudes of each $B_{(s)} \rightarrow D_{(s)}P$ channels are then

$$A(B^- \rightarrow D^0 \pi^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* (\xi_{ext} + \mathcal{M}_{ext} + \xi_{int} + \mathcal{M}_{int}) , \quad (47)$$

$$A(B^- \rightarrow D^0 K^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* (\xi_{ext} + \mathcal{M}_{ext} + \xi_{int} + \mathcal{M}_{int}) , \quad (48)$$

$$A(\bar{B}^0 \rightarrow D^+ \pi^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* (\xi_{ext} + \mathcal{M}_{ext} + \xi_{exc} + \mathcal{M}_{exc}) , \quad (49)$$

$$A(\bar{B}^0 \rightarrow D^+ K^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* (\xi_{ext} + \mathcal{M}_{ext}) , \quad (50)$$

$$A(\bar{B}^0 \rightarrow D_s^+ K^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* (\xi_{exc} + \mathcal{M}_{exc}) , \quad (51)$$

$$A(\bar{B}^0 \rightarrow D^0 \pi^0) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* \frac{1}{\sqrt{2}} (-(\xi_{int} + \mathcal{M}_{int}) + (\xi_{exc} + \mathcal{M}_{exc})) , \quad (52)$$

$$A(\bar{B}^0 \rightarrow D^0 \bar{K}^0) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* (\xi_{int} + \mathcal{M}_{int}) , \quad (53)$$

$$A(\bar{B}^0 \rightarrow D^0 \eta_n) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* \frac{1}{\sqrt{2}} (\xi_{int} + \mathcal{M}_{int} + \xi_{exc} + \mathcal{M}_{exc}) , \quad (54)$$

$$A(\bar{B}_s^0 \rightarrow D^+ \pi^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* (\xi_{exc} + \mathcal{M}_{exc}) , \quad (55)$$

$$A(\bar{B}_s^0 \rightarrow D_s^+ \pi^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* (\xi_{ext} + \mathcal{M}_{ext}) , \quad (56)$$

$$A(\bar{B}_s^0 \rightarrow D_s^+ K^-) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* (\xi_{ext} + \mathcal{M}_{ext} + \xi_{exc} + \mathcal{M}_{exc}) , \quad (57)$$

$$A(\bar{B}_s^0 \rightarrow D^0 \pi^0) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* \frac{1}{\sqrt{2}} (\xi_{exc} + \mathcal{M}_{exc}) , \quad (58)$$

$$A(\bar{B}_s^0 \rightarrow D^0 \bar{K}^0) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* (\xi_{int} + \mathcal{M}_{int}) , \quad (59)$$

$$A(\bar{B}_s^0 \rightarrow D^0 \eta_n) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* \frac{1}{\sqrt{2}} (\xi_{exc} + \mathcal{M}_{exc}) , \quad (60)$$

$$A(\bar{B}_s^0 \rightarrow D^0 \eta_s) = \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* (\xi_{int} + \mathcal{M}_{int}) . \quad (61)$$

It should be noticed that, in (54), (60) and (61), the decay amplitudes are for the mixing basis of η and η' . For the physical state η and η' , the decay amplitudes are

$$A(\bar{B}^0 \rightarrow D^0 \eta) = A(\bar{B}^0 \rightarrow D^0 \eta_n) \cos \phi , \quad (62)$$

$$A(\bar{B}^0 \rightarrow D^0 \eta') = A(\bar{B}^0 \rightarrow D^0 \eta_n) \sin \phi , \quad (63)$$

$$A(\bar{B}_s^0 \rightarrow D^0 \eta) = A(\bar{B}_s^0 \rightarrow D^0 \eta_n) \cos \phi - A(\bar{B}_s^0 \rightarrow D^0 \eta_s) \sin \phi , \quad (64)$$

$$A(\bar{B}_s^0 \rightarrow D^0 \eta') = A(\bar{B}_s^0 \rightarrow D^0 \eta_n) \sin \phi + A(\bar{B}_s^0 \rightarrow D^0 \eta_s) \cos \phi . \quad (65)$$

B. Amplitudes for $B_{(s)} \rightarrow D_{(s)}V$ and $B_{(s)} \rightarrow D_{(s)}^*P$ decays

For the channels of $B_{(s)} \rightarrow D_{(s)}V$ and $D_{(s)}^*P$, no transverse polarization of the vector mesons will contribute. In the leading power contribution, the formulae of $B \rightarrow DV$ and $B \rightarrow D^*P$ are the same as that of $B_{(s)} \rightarrow D_{(s)}P$ decays, except some substitutions.

For $B_{(s)} \rightarrow D_{(s)}V$, the following substitutions should be done for the formula ξ_i and $\mathcal{M}_i$:

$$\phi_P \rightarrow \phi_V, \phi_P^p \rightarrow -\phi_V^s, \phi_P^t \rightarrow -\phi_V^t, m_0 \rightarrow m_V, f_P \rightarrow f_V. \quad (66)$$

ϕ_V , ϕ_V^s and ϕ_V^t are the light cone distribution amplitudes of vector mesons, which we defined before. And m_V , f_V are the mass and the decay constant of the vector meson related.

For $B_{(s)} \rightarrow D_{(s)}^*P$, the substitutions in the formula ξ_i and $\mathcal{M}_i$ should be done as:

$$m_D \rightarrow m_{D^*}, f_D \rightarrow f_{D^*}, \phi_D(x_2) \rightarrow \phi_{D^*}(x_2). \quad (67)$$

Making the following substitutions in equations (47)-(65), we can get the final decay amplitude for each $B \rightarrow D^*P$ decays:

$$D^+ \rightarrow D^{*+}, D^0 \rightarrow D^{*0}, D_s^+ \rightarrow D_s^{*+}. \quad (68)$$

And the formulae for $B \rightarrow DV$ can be obtained through the substitutions

$$\pi \rightarrow \rho, K \rightarrow K^*, \eta_n \rightarrow \omega, \eta_s \rightarrow \phi \quad (69)$$

in equations (47)-(61).

C. Amplitudes for $B_{(s)} \rightarrow D_{(s)}^*V$ decays

For $B \rightarrow D_{(s)}^*V$ decays, both longitudinal and transverse polarization can contribute. For the longitudinal polarization, the amplitudes can be obtained by carrying out the substitutions referred in (66) and (67), when only the leading power contribution is taken into consideration. The transverse polarized contribution is suppressed by r or r_V , $r_V \equiv m_V/m_B$. Although the transverse polarization will not give the leading power contribution, we still list the analytic formulae for

transverse polarization ξ_{ext}^T , ξ_{int}^T and ξ_{exc}^T

$$\begin{aligned}\xi_{\text{ext}}^T &= 8\pi C_F m_B^4 f_V \int_0^1 dx_1 dx_2 \int_0^{1/\Lambda} b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_D^T(x_2) \\ &\quad \times r_V \left[E_e(t_e^{(1)}) h(x_1, x_2, b_1, b_2) S_t(x_2) \left(-\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} (1 + 2r) \right) \right. \\ &\quad \left. + E_e(t_e^{(2)}) h(x_2, x_1, b_2, b_1) S_t(x_1) r(r+1) \left(-\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} \right) \right],\end{aligned}\quad (70)$$

$$\begin{aligned}\xi_{\text{int}}^T &= 8\pi C_F m_B^4 f_{D^*} \int_0^1 dx_1 dx_3 \int_0^{1/\Lambda} b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) \\ &\quad \times r \left\{ \left[\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} \left(-\phi_V^t(x_3) - r_V((x_3 - 3)\phi_V^a(x_3) + (x_3 - 1)\phi_V^v(x_3)) \right) \right. \right. \\ &\quad \left. \left. - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} \left(\phi_V^t(x_3) - r_V((x_3 - 1)\phi_V^a(x_3) - (x_3 - 3)\phi_V^v(x_3)) \right) \right] \right. \\ &\quad \times E_i(t_i^{(1)}) h(x_1, x_3(1 - r^2), b_1, b_3) S_t(x_3) \\ &\quad \left. + r_V \left[\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} (\phi_V^a(x_3) - \phi_V^v(x_3)) + i \epsilon_D^{*T} \cdot \epsilon_V^{*T} (\phi_V^a(x_3) - \phi_V^v(x_3)) \right] \right. \\ &\quad \left. \times E_i(t_i^{(2)}) h(x_3, x_1(1 - r^2), b_3, b_1) S_t(x_1) \right\},\end{aligned}\quad (71)$$

$$\begin{aligned}\xi_{\text{exc}}^T &= 8\pi C_F m_B^4 f_B \int_0^1 dx_2 dx_3 \int_0^{1/\Lambda} b_2 db_2 b_3 db_3 \phi_D^T(x_2) \\ &\quad \times \left[E_a(t_a^{(1)}) h_a(x_2, x_3(1 - r^2), b_2, b_3) S_t(x_3) \right. \\ &\quad \times \left(\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} [r^2 \phi_V^t(x_3) - r r_V(x_3 + 1) \phi_V^a(x_3) + r r_V(1 - x_3) \phi_V^v(x_3)] \right. \\ &\quad \left. + i \epsilon_D^{*T} \cdot \epsilon_V^{*T} [-r^2 \phi_V^t(x_3) + r r_V(x_3 - 1) \phi_V^a(x_3) + r r_V(x_3 + 1) \phi_V^v(x_3)] \right) \\ &\quad + E_a(t_a^{(2)}) h_a(x_3, x_2(1 - r^2), b_3, b_2) S_t(x_2) r r_V \\ &\quad \times \left(\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} ((1 + x_2) \phi_V^a(x_3) + (1 - x_2) \phi_V^v(x_3)) \right. \\ &\quad \left. - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} ((1 - x_2) \phi_V^a(x_3) + (1 + x_2) \phi_V^v(x_3)) \right) \left. \right].\end{aligned}\quad (72)$$

The evolution factors in the amplitudes are the same as those in (35) and (36) after substituting $S_V(t)$ for $S_P(t)$. For the nonfactorizable amplitudes, the factorization formulas involve the

kinematic variables of all the three mesons. Their expressions are

$$\begin{aligned}
\mathcal{M}_{\text{ext}}^T = & 16\pi\sqrt{2N_c}C_Fm_B^4 \int_0^1 [dx] \int_0^{1/\Lambda} b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) \phi_D^T(x_2) \\
& \times r_V \left[E_b(t_b^{(1)}) h_b^{(1)}(x_i, b_i) \right. \\
& \times \left(\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} x_3 (\phi_V^a(x_3) - \phi_V^v(x_3)) + i \epsilon_D^{*T} \cdot \epsilon_V^{*T} x_3 (\phi_V^a(x_3) - \phi_V^v(x_3)) \right) \\
& + E_b(t_b^{(2)}) h_b^{(2)}(x_i, b_i) \\
& \times \left\{ \epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} ((1-x_3)(1-2r)\phi_V^a(x_3) + (x_3-1)\phi_V^v(x_3)) \right. \\
& \left. \left. - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} ((x_3-1)\phi_V^a(x_3) + (1-2r)(1-x_3)\phi_V^v(x_3)) \right\} \right], \tag{73}
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{\text{int}}^T = & 16\pi\sqrt{2N_c}C_Fm_B^4 \int_0^1 [dx] \int_0^{1/\Lambda} b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_D^T(x_2) \\
& \times r \left[E_d(t_d^{(2)}) h_d^{(2)}(x_i, b_i) \right. \\
& \times \left(\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} ((x_2-1)\phi_V^t(x_3) + r_V(\phi_V^a(x_3) + \phi_V^v(x_3))) \right. \\
& \left. - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} ((1-x_2)\phi_V^t(x_3) + r_V(\phi_V^a(x_3) + \phi_V^v(x_3))) \right) \\
& + E_d(t_d^{(1)}) h_d^{(1)}(x_i, b_i) \\
& \times \left(\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} (2r_V(x_3-1)\phi_V^a(x_3) - x_2\phi_V^t(x_3)) \right. \\
& \left. \left. - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} (x_2\phi_V^t(x_3) + 2r_V(x_3-1)\phi_V^v(x_3)) \right) \right], \tag{74}
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{\text{exc}}^T = & 16\pi\sqrt{2N_c}C_Fm_B^4 \int_0^1 [dx] \int_0^{1/\Lambda} b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_D^T(x_2) \\
& \times \left\{ E_f(t_f^{(1)}) h_f^{(1)}(x_i, b_i) \phi_V^t(x_3) \right. \\
& \times \left(\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} (x_2 r^2 - r_V^2 x_3) - i \epsilon_D^{*T} \cdot \epsilon_V^{*T} (x_2 r^2 + r_V^2 x_3) \right) \\
& + E_f(t_f^{(2)}) h_f^{(2)}(x_i, b_i) \\
& \times \left[\epsilon^{n\bar{n}} \epsilon_D^{*T} \epsilon_V^{*T} (-2r r_V \phi_V^a(x_3) - r^2(x_2-1)\phi_V^t(x_3) + r_V^2(x_3-1)\phi_V^t(x_3)) \right. \\
& \left. \left. + i \epsilon_D^{*T} \cdot \epsilon_V^{*T} (((x_2-1)r^2 + r_V^2(x_2-1))\phi_V^t(x_3) + 2r r_V \phi_V^v(x_3)) \right] \right\}. \tag{75}
\end{aligned}$$

The h 's and $h^{(j)}$ functions in the amplitudes here are the same as defined in (35,36,42-44).

If carrying out the substitutions in (68) and (69) together, we can get the final decay amplitude of each polarization for $B \rightarrow D^*V$ decays.

TABLE II: The smallest χ^2 for each kind of the D meson DAs

	$\Phi^{(Gen)}$	$\Phi^{(MGen)}$	$\Phi^{(KLS)}$	$\Phi^{(GN)}$	$\Phi^{(KKQT)}$	$\Phi^{(Huang)}$
χ_{min}^2	34.1	33.6	156.9	112.3	45.0	41.9

IV. NUMERICAL CALCULATIONS AND DISCUSSIONS

In this section the numerical results of our calculation will be given. The parameters of $D_{(s)}^{(*)}$ meson we use are

$$\begin{aligned}
m_{D^-} &= 1.869\text{GeV}, & m_{D_s^-} &= 1.968\text{GeV}, \\
m_{D^{*-}} &= 2.010\text{GeV}, & m_{D_s^{*-}} &= 2.112\text{GeV}, \\
f_{D^-} &= 223\text{MeV}, & f_{D_s^-} &= 274\text{MeV}.
\end{aligned} \tag{76}$$

A lot of study has been made on the decay constants of $D_{(s)}$ mesons. Here we take the values from ref. [20]. Since there are no experimental results of the decay constants of $D_{(s)}^*$ mesons, we use the relations between f_D and f_{D^*} derived from HQET:

$$f_{D^{*-}} = \sqrt{\frac{m_{D^-}}{m_{D^{*-}}}} f_{D^-}, \quad f_{D_s^{*-}} = \sqrt{\frac{m_{D_s^-}}{m_{D_s^{*-}}}} f_{D_s^-}, \tag{77}$$

which is different from ref.[21].

With the D meson wave functions we get, we can calculate the amplitudes easily. And the decay width can be got by

$$\Gamma = \frac{1}{32\pi} G_F^2 |V_{cb}|^2 |V_{ud}|^2 m_B^7 (1 - r^2) |A|^2, \tag{78}$$

with A the decay amplitude defined in eqs.(47-61) The branch ratio is

$$Br = \Gamma \hbar / \tau_{B_{(s)}}, \tag{79}$$

with $\tau_{B_{(s)}}$ as the life time of $B_{(s)}$ meson. We take $\tau_{B^-} = 1.674 \times 10^{-12} s$, $\tau_{\bar{B}^0} = 1.542 \times 10^{-12} s$, $\tau_{\bar{B}_s^0} = 1.466 \times 10^{-12} s$, and $G_F = 1.16639 \times 10^{-5}$.

A. results of fitting

Since the $B \rightarrow DP$ decay channels have been measured in high precision, we use these experimental results to fit out the parameters of the candidate D meson DAs. Here we don't use




FIG. 4: The D meson distribution amplitude $\phi_D^{MGen}(0.8, 0)$ (the blue, solid line) and D_s meson distribution amplitude $\phi_{D_s}^{MGen}(0.5, 0)$ (the red, dotted line)

the experimental results containing η or η' in the final states because there are uncertainties from the mixing. The six decay channels we used to fit out the D meson wave function parameters are $B^- \rightarrow D^0 \pi^-$, $B^- \rightarrow D^0 K^-$, $\bar{B}^0 \rightarrow D^+ \pi^-$, $\bar{B}^0 \rightarrow D^+ K^-$, $\bar{B}^0 \rightarrow D^0 \pi^0$, $\bar{B}^0 \rightarrow D^0 \bar{K}^0$. The experimental results of these channels are listed in table III, which are from ref.[22]. The formula we used for fitting is

$$\chi^2 = \sum_i \frac{(Br_i^{ex} - Br_i^{th})^2}{\sigma_i^2}. \quad (80)$$

The i means the summation over the six decay channels. $Br_i^{ex}(Br_i^{th})$ is the experimental (theoretical) value of branch ratio, and σ_i is the uncertainty of the experimental value. In table II we list the smallest χ^2 we get for all the D meson DAs. Easy to see, except $\Phi^{(KLS)}$ and $\Phi^{(GN)}$, all the other DAs have a small χ_{min}^2 . The $\Phi^{(MGen)}$ is the best one, with its parameters fixed as $C_D = 0.8$, $\omega = 0.1$. We will use this D meson wave function for the following numerical calculations of all the decay channels. For the D_s meson, we use $C_D = 0.5$, $\omega = 0.2$, with a little SU(3) breaking effect. In this case, we can see from Fig. 4 that the $\bar{s}$ quark in D_s meson has a little larger momentum fraction than the $\bar{d}$ quark in the D meson, which characterize the little larger mass of s quark.

Because the mass difference between the vector meson $D_{(s)}^*$ and pseudoscalar meson $D_{(s)}$ is small, so we adopt the same DA for them.

TABLE III: Branching ratios of $B_{(s)} \rightarrow DP$ decays calculated in pQCD approach with experimental data (in units of 10^{-4})

	Experimental results	Our results
$B^- \rightarrow D^0 \pi^-$	47.5 ± 1.9	$51.1^{+29.5+4.3+1.5}_{-20.7-7.5-1.5}$
$B^- \rightarrow D^0 K^-$	3.83 ± 0.45	$4.00^{+2.35+0.63+0.12}_{-1.64-0.93-0.12}$
$\bar{B}^0 \rightarrow D^+ \pi^-$	26.5 ± 1.5	$26.9^{+17.8+5.5+0.8}_{-11.7-7.3-0.8}$
$\bar{B}^0 \rightarrow D^+ K^-$	2.04 ± 0.57	$2.43^{+1.56+0.63+0.07}_{-1.01-0.71-0.07}$
$\bar{B}^0 \rightarrow D^0 \pi^0$	2.61 ± 0.25	$1.98^{+0.67+0.51+0.06}_{-0.66-0.63-0.06}$
$\bar{B}^0 \rightarrow D^0 \bar{K}^0$	0.523 ± 0.066	$0.22^{+0.08+0.06+0.01}_{-0.07-0.06-0.01}$
$\bar{B}^0 \rightarrow D^0 \eta$	2.02 ± 0.21	$2.46^{+0.97+0.30+0.07}_{-0.77-0.30-0.07}$
$\bar{B}^0 \rightarrow D^0 \eta'$	1.26 ± 0.21	$1.65^{+0.65+0.20+0.05}_{-0.52-0.20-0.05}$
$\bar{B}^0 \rightarrow D_s^+ K^-$	0.269 ± 0.054	$0.48^{+0.16+0.11+0.01}_{-0.12-0.11-0.01}$
$\bar{B}_s^0 \rightarrow D^+ \pi^-$		$2.32^{+0.82+0.32+0.07}_{-0.61-0.39-0.07} \times 10^{-2}$
$\bar{B}_s^0 \rightarrow D^0 \pi^0$		$1.15^{+0.36+0.19+0.04}_{-0.29-0.20-0.04} \times 10^{-2}$
$\bar{B}_s^0 \rightarrow D^0 \bar{K}^0$		$3.96^{+1.49+0.88+0.11}_{-1.25-0.99-0.12}$
$\bar{B}_s^0 \rightarrow D^0 \eta$		$0.14^{+0.05+0.03+0.00}_{-0.05-0.03-0.00}$
$\bar{B}_s^0 \rightarrow D^0 \eta'$		$0.32^{+0.11+0.03+0.01}_{-0.10-0.04-0.01}$
$\bar{B}_s^0 \rightarrow D_s^+ \pi^-$	$38 \pm 3 \pm 13$	$21.3^{+11.4+6.9+0.6}_{-8.1-6.8-0.6}$
$\bar{B}_s^0 \rightarrow D_s^+ K^-$		$1.71^{+0.92+0.58+0.05}_{-0.65-0.55-0.05}$

B. Results for all the related channels and discussions

Our numerical results are listed in table III, table IV, table V and table VI. The first error in these entries are caused by the hadronic parameters in $B_{(s)}$ meson wave function (the decay constant and the shape parameter). We take $f_B = 0.19 \pm 0.025$, $\omega_b = 0.40 \pm 0.05$ and $f_{B_s} = 0.24 \pm 0.03$, $\omega_b^{B_s} = 0.50 \pm 0.05$. The second error arise from the higher order perturbative QCD corrections: the choice of the hard scales, defined in (37) and (46), which vary from $0.75t$ to $1.25t$, and the uncertainty of $\Lambda_{QCD}^{(4)} = 0.25 \pm 0.05$. The third error is from the uncertainties of the CKM matrix elements. In our calculation, we use

$$V_{cb} = (41.61^{+0.62}_{-0.63}) \times 10^{-3}, \quad V_{ud} = 0.97385^{+0.00024}_{-0.00023}, \quad V_{us} = 0.22715^{+0.00101}_{-0.00100}. \quad (81)$$

Among them, the hadronic inputs always gives rise to the largest uncertainty, and the CKM matrix elements contribute little.

The first six channels in table III are input values of the χ^2 fit program. Although we get a

TABLE IV: Branching ratios of $B_{(s)} \rightarrow DV$ decays calculated in pQCD approach with experimental data (in units of 10^{-4})

	Experimental results	Our results
$B^- \rightarrow D^0 \rho^-$	134 ± 18	$113^{+68.1+18.8+3.3}_{-45.9-26.1-3.3}$
$B^- \rightarrow D^0 K^{*-}$	5.29 ± 0.45	$6.49^{+3.86+0.12+0.20}_{-2.68-1.58-0.20}$
$\bar{B}^0 \rightarrow D^+ \rho^-$	75 ± 12	$69.6^{+45.0+13.1+2.0}_{-29.7-17.8-2.0}$
$\bar{B}^0 \rightarrow D^+ K^{*-}$	4.60 ± 0.78	$4.07^{+2.61+0.94+0.12}_{-1.69-1.11-0.12}$
$\bar{B}^0 \rightarrow D^0 \rho^0$	2.91 ± 0.40	$1.79^{+0.65+0.46+0.05}_{-0.62-0.59-0.05}$
$\bar{B}^0 \rightarrow D^0 \omega$	2.60 ± 0.29	$4.25^{+1.58+0.85+0.12}_{-1.21-0.73-0.12}$
$\bar{B}^0 \rightarrow D^0 \bar{K}^{*0}$	0.423 ± 0.064	$0.26^{+0.10+0.08+0.01}_{-0.08-0.07-0.01}$
$\bar{B}^0 \rightarrow D_s^+ K^{*-}$	< 8	$1.94^{+0.66+0.41+0.06}_{-0.47-0.38-0.06}$
$\bar{B}_s^0 \rightarrow D^+ \rho^-$		$0.11^{+0.03+0.02+0.00}_{-0.03-0.02-0.00}$
$\bar{B}_s^0 \rightarrow D^0 \rho^0$		$(5.42^{+1.57+0.96+0.16}_{-1.39-1.11-0.16}) \times 10^{-2}$
$\bar{B}_s^0 \rightarrow D^0 \bar{K}^{*0}$		$4.52^{+1.70+1.15+0.13}_{-1.43-1.30-0.13}$
$\bar{B}_s^0 \rightarrow D^0 \omega$		$(4.71^{+1.35+0.88+0.14}_{-1.18-0.92-0.14}) \times 10^{-2}$
$\bar{B}_s^0 \rightarrow D^0 \phi$		$0.30^{+0.11+0.07+0.01}_{-0.10-0.08-0.01}$
$\bar{B}_s^0 \rightarrow D_s^+ \rho^-$		$51.0^{+26.9+16.6+1.48}_{-19.2-16.2-1.48}$
$\bar{B}_s^0 \rightarrow D_s^+ K^{*-}$		$3.02^{+1.62+0.88+0.10}_{-1.16-0.91-0.10}$

reasonable χ^2 in the fit, the branching ratio of $\bar{B}^0 \rightarrow D^0 \bar{K}^0$ is about only half of the experimental value. Comparing with the color suppressed diagrams, the annihilation diagrams contributes little for $\bar{B}^0 \rightarrow D^0 \bar{K}^0$ and $\bar{B}^0 \rightarrow D^0 \pi^0$. So we can take them for a compare. They have different CKM elements ($V_{cb}V_{us}^*$ for the former and $V_{cb}V_{ud}^*$ for the latter). Taking the factor $\frac{1}{\sqrt{2}}$ in flavor wave function of π meson into account, the $\text{Br}(\bar{B}^0 \rightarrow D^0 \bar{K}^0)$ is roughly one tenth of $\text{Br}(\bar{B}^0 \rightarrow D^0 \pi^0)$. So this value of $\text{Br}(\bar{B}^0 \rightarrow D^0 \bar{K}^0)$ is theoretically reasonable. The similar argument is valid for $\text{Br}(\bar{B}^0 \rightarrow D^{*0} \bar{K}^0)$.

Although we use only six $B \rightarrow DP$ channels to fix the D meson wave function, the results of most other channels especially those $B \rightarrow DV$ and D^*P channels agree very well with the current experimental measurements. Easy to find that, $\text{Br}(B \rightarrow D\omega)$ is twice larger than $\text{Br}(B \rightarrow D\rho)$, while their experimental results are near to each other. Both these two channels receive contributions from the color suppressed diagrams and annihilation diagrams. They are at the same order magnitude for these two processes. For color suppressed diagrams, the $d\bar{d}$ of the flavor part contributes. While the $u\bar{u}$ part contributes to the annihilation diagrams. Amplitudes of these

TABLE V: Branching ratios of $B_{(s)} \rightarrow D^* P$ decays calculated in pQCD approach with experimental data (in units of 10^{-4})

	Experimental results	Our results
$B^- \rightarrow D^{*0} \pi^-$	52.8 ± 2.8	$50.4^{+29.4+4.4+1.5}_{-20.4-7.3-1.5}$
$B^- \rightarrow D^{*0} K^-$	3.6 ± 1.0	$3.98^{+2.33+0.62+0.12}_{-1.62-0.92-0.12}$
$\bar{B}^0 \rightarrow D^{*+} \pi^-$	26.2 ± 1.3	$26.0^{+17.3+5.34+0.75}_{-11.4-7.09-0.76}$
$\bar{B}^0 \rightarrow D^{*+} K^-$	2.04 ± 0.47	$2.37^{+1.52+0.62+0.07}_{-0.99-0.69-0.07}$
$\bar{B}^0 \rightarrow D^{*0} \pi^0$	1.71 ± 0.28	$2.11^{+0.76+0.52+0.06}_{-0.69-0.62-0.06}$
$\bar{B}^0 \rightarrow D^{*0} \eta$	1.80 ± 0.31	$2.60^{+1.01+0.29+0.08}_{-0.83-0.33-0.08}$
$\bar{B}^0 \rightarrow D^{*0} \eta'$	1.21 ± 0.40	$1.74^{+0.68+0.19+0.05}_{-0.55-0.22-0.05}$
$\bar{B}^0 \rightarrow D^{*0} \bar{K}^0$	0.36 ± 0.12	$0.24^{+0.09+0.06+0.01}_{-0.07-0.06-0.01}$
$\bar{B}^0 \rightarrow D_s^{*+} K^-$	0.200 ± 0.064	$0.47^{+0.16+0.12+0.01}_{-0.12-0.11-0.01}$
$\bar{B}_s^0 \rightarrow D^{*+} \pi^-$		$(2.27^{+0.81+0.31+0.01}_{-0.60-0.38-0.01}) \times 10^{-2}$
$\bar{B}_s^0 \rightarrow D^{*0} \pi^0$		$(1.13^{+0.35+0.18+0.03}_{-0.28-0.19-0.03}) \times 10^{-2}$
$\bar{B}_s^0 \rightarrow D^{*0} \bar{K}^0$		$4.27^{+1.54+0.88+0.12}_{-1.35-1.05-0.12}$
$\bar{B}_s^0 \rightarrow D^{*0} \eta$		$0.15^{+0.06+0.03+0.00}_{-0.05-0.03-0.00}$
$\bar{B}_s^0 \rightarrow D^{*0} \eta'$		$0.33^{+0.12+0.03+0.01}_{-0.10-0.03-0.01}$
$\bar{B}_s^0 \rightarrow D_s^{*+} \pi^-$		$24.2^{+11.2+7.8+0.7}_{-7.2-7.7-0.7}$
$\bar{B}_s^0 \rightarrow D_s^{*+} K^-$		$1.65^{+0.90+0.56+0.05}_{-0.63-0.53-0.05}$

two kinds of diagrams have the same sign in the $B \rightarrow D\omega$ decay but different sign in the $B \rightarrow D\rho$ decay due to isospin. Similar situation exists for $\text{Br}(B \rightarrow D^*\omega)$ and $\text{Br}(B \rightarrow D^*\rho)$.

The $\bar{B}^0 \rightarrow D_s^+ K^-$ decay is a kind of pure annihilation type decays dominant by W exchange diagram. Our result is larger than the experiments and also larger than the previous pQCD calculations [23] due to the change of D meson wave functions. The annihilation type diagrams are power suppressed in pQCD approach, which is more sensitive to the hadronic wave functions.

For the decays $B \rightarrow D_{(s)}^* V$ in table VI, we also estimate the ratios of transverse polarized contribution $R_T = |A_T|^2 / (|A_T|^2 + |A_L|^2)$. We should mention that, these results are just indicative, because transverse polarizations are power suppressed by r_V or r to make it more sensitive to small parameters and higher order contributions than the longitudinal contribution. Although the transverse polarization is suppressed in $B \rightarrow D^* V$ decays, in some channels, such as $B \rightarrow D^{*0} \rho^0$ and $B \rightarrow D^{*0} \bar{K}^{*0}$, etc, it has 40% contributions. The reason is that the dominant contribution in these channels is from $\mathcal{M}_{int}$ in eq.(39), which is x_3 suppressed, while the transverse contribution

TABLE VI: Predicted branching ratios of $B_{(s)} \rightarrow D^*V$ decays with experimental data (in units of 10^{-4}) together with the percentage of transverse polarizations R_T

Experimental BRs			BRs in pQCD	R_T
$B^- \rightarrow D^{*0}\rho^-$			$117^{+71.0+19.9+3.4}_{-48.0-27.0-3.4}$	0.04
$B^- \rightarrow D^{*0}K^{*-}$	8.3 ± 1.5	0.14	$6.82^{+4.14+1.22+0.21}_{-2.80-1.65-0.21}$	0.06
$\bar{B}^0 \rightarrow D^{*+}\rho^-$			$79.4^{+52.6+15.9+2.3}_{-34.5-21.2-2.3}$	0.15
$\bar{B}^0 \rightarrow D^{*+}K^{*-}$	3.20 ± 0.67		$4.88^{+3.18+1.16+0.15}_{-2.08-1.41-0.15}$	0.19
$\bar{B}^0 \rightarrow D^{*0}\rho^0$	3.73 ± 0.99		$3.69^{+1.52+0.57+0.11}_{-1.23-0.63-0.11}$	0.47
$\bar{B}^0 \rightarrow D^{*0}\omega$	2.68 ± 0.50		$5.83^{+2.14+0.84+0.17}_{-1.84-1.10-0.17}$	0.24
$\bar{B}^0 \rightarrow D^{*0}\bar{K}^{*0}$	< 0.69		$0.51^{+0.20+0.08+0.02}_{-0.17-0.10-0.02}$	0.44
$\bar{B}^0 \rightarrow D_s^{*+}K^{*-}$			$1.97^{+0.58+0.37+0.06}_{-0.49-0.43-0.06}$	0.02
$\bar{B}_s^0 \rightarrow D^{*+}\rho^-$			$0.11^{+0.03+0.02+0.00}_{-0.03-0.02-0.00}$	0.01
$\bar{B}_s^0 \rightarrow D^{*0}\rho^0$			$(5.35^{+1.44+0.96+0.16}_{-1.39-1.05-0.16}) \times 10^{-2}$	0.01
$\bar{B}_s^0 \rightarrow D^{*0}\bar{K}^{*0}$			$8.43^{+3.30+1.68+0.24}_{-2.67-1.83-0.25}$	0.43
$\bar{B}_s^0 \rightarrow D^{*0}\omega$			$(4.60^{+1.50+0.89+0.14}_{-1.11-0.84-0.14}) \times 10^{-2}$	0.01
$\bar{B}_s^0 \rightarrow D^{*0}\phi$			$0.51^{+0.19+0.09+0.02}_{-0.17-0.11-0.02}$	0.37
$\bar{B}_s^0 \rightarrow D_s^{*+}\rho^-$			$56.9^{+30.4+19.1+1.65}_{-21.6-18.3-1.66}$	0.13
$\bar{B}_s^0 \rightarrow D_s^{*+}K^{*-}$			$3.47^{+1.96+1.07+0.11}_{-1.35-1.06-0.11}$	0.17

in eq.(74) is only r suppressed. They are comparable numerically to make a large contribution for transverse polarizations in these color suppressed channels. This mechanism is different from those charmless B decays where the dominant transverse polarizations are from the space like penguin (penguin annihilation) contributions [27]. Here the annihilation type contributions are mainly from W exchange diagrams contributing little to transverse polarizations.

For those previous calculated channels in pQCD approach [7, 9], our results are slightly different due to parameter changes. Most of the $B^0(B^\pm)$ decay channels are measured by the two B factories, which are consistent with our calculations. For the B_s decays, only one channel is measured. Our predictions will soon be tested by the LHCb experiments.

For comparison with other methods, we also give the form factors at the maximal recoil

$$\xi_+^{B \rightarrow D} = 0.52^{+0.15+0.05}_{-0.12-0.07}, \quad \xi_+^{B_s \rightarrow D_s} = 0.46^{+0.11+0.07}_{-0.09-0.08}. \quad (82)$$

These are comparable with other methods [28].

If applying the naive factorization approach, we can get

$$A(\bar{B}^0 \rightarrow D^+\pi^-) = i\frac{G_F}{\sqrt{2}}V_{cb}V_{ud}^*(M_B^2 - M_D^2)f_\pi F^{B \rightarrow D}(M_\pi^2)a_1(D\pi), \quad (83)$$

$$\sqrt{2}A(\bar{B}^0 \rightarrow D^0\pi^0) = -i\frac{G_F}{\sqrt{2}}V_{cb}V_{ud}^*(M_B^2 - M_\pi^2)f_DF^{B \rightarrow \pi}(M_D^2)a_2(D\pi). \quad (84)$$

Substituting our results for $A(\bar{B}^0 \rightarrow D^+\pi^-)$ and $A(\bar{B}^0 \rightarrow D^0\pi^0)$ in eq.(83) and (84), we can extract the BSW parameters a_1 and a_2 from our pQCD approach

$$|a_2/a_1| = 0.49, \text{ Arg}(a_2/a_1) = -37.6^\circ. \quad (85)$$

If the annihilation diagrams' contribution is excluded, the results are

$$|a_2/a_1| = 0.54, \text{ Arg}(a_2/a_1) = -59.0^\circ. \quad (86)$$

Indeed, the large $|a_2/a_1|$ implies that the color suppressed decays not very suppressed as previous expected [3]. The relative strong phase between the two contributions is not small as naive expectations. These results are consistent with recent direct studies from experiments [24]. But the difference is that our results come from direct dynamical calculation not from fit.

V. CONCLUSION

In this paper, we calculate the branch ratios of $B_{(s)} \rightarrow D_{(s)}P$, $D_{(s)}^*P$, $D_{(s)}V$ and $D_{(s)}^*V$ channels, with the D meson wave function obtained through fitting. We also calculate the ratios of transverse polarized contributions in $B \rightarrow D^*V$ decays. Most of the results agree well with the experiments. It seems that there's a disagreement with the experimental data in the relative size of branching ratios for $B \rightarrow D^{(*)}\rho$ and $B \rightarrow D^{(*)}\omega$. Some channels of the $B \rightarrow D^*V$ decays may receive large contribution from the transverse polarization. The results of $\bar{B}_s^0 \rightarrow D_{(s)}P$, $D_{(s)}V$, $D_{(s)}^*P$, $D_{(s)}^*V$ decays will be tested in the future experiments.

Acknowledgments

We thank Wei Wang and Yu-Ming Wang a lot, for the fruitful discussions. This work is partly supported by National Science Foundation of China under the Grant Numbers 10735080, 10625525 and 10525523.

APPENDIX A: PQCD FUNCTIONS

Jet function appears in (31)-(33), (70)-(72) is

$$S_t(x) = \frac{2^{1+2c} \Gamma(3/2 + c)}{\sqrt{\pi} \Gamma(1 + c)} [x(1 - x)]^c \quad (\text{A1})$$

the value of c in the above equation is 0.5 in this paper. And the $S_j(x_i)$ ($j = B, C, P$ or V) functions in Sudakov form factors in (34) and (41) are

$$S_B(t) = s\left(x_1 \frac{m_B}{\sqrt{2}}, b_1\right) + 2 \int_{1/b_1}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})), \quad (\text{A2})$$

$$S_C(t) = s\left(x_2 \frac{m_B}{\sqrt{2}}, b_2\right) + 2 \int_{1/b_2}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})), \quad (\text{A3})$$

$$S_V(t) = S_P(t) = s\left(x_3 \frac{m_B}{\sqrt{2}}, b_3\right) + s\left((1 - x_3) \frac{m_B}{\sqrt{2}}, b_3\right) + 2 \int_{1/b_3}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})), \quad (\text{A4})$$

with the quark anomalous dimension $\gamma_q = -\alpha_s/\pi$. The explicit form for the function $s(Q, b)$ is:

$$\begin{aligned} s(Q, b) = & \frac{A^{(1)}}{2\beta_1} \hat{q} \ln\left(\frac{\hat{q}}{\hat{b}}\right) - \frac{A^{(1)}}{2\beta_1} (\hat{q} - \hat{b}) + \frac{A^{(2)}}{4\beta_1^2} \left(\frac{\hat{q}}{\hat{b}} - 1\right) - \left[\frac{A^{(2)}}{4\beta_1^2} - \frac{A^{(1)}}{4\beta_1} \ln\left(\frac{e^{2\gamma_E - 1}}{2}\right)\right] \ln\left(\frac{\hat{q}}{\hat{b}}\right) \\ & + \frac{A^{(1)}\beta_2}{4\beta_1^3} \hat{q} \left[\frac{\ln(2\hat{q}) + 1}{\hat{q}} - \frac{\ln(2\hat{b}) + 1}{\hat{b}}\right] + \frac{A^{(1)}\beta_2}{8\beta_1^3} [\ln^2(2\hat{q}) - \ln^2(2\hat{b})], \end{aligned} \quad (\text{A5})$$

where the variables are defined by

$$\hat{q} \equiv \ln[Q/(\sqrt{2}\Lambda)], \quad \hat{b} \equiv \ln[1/(b\Lambda)], \quad (\text{A6})$$

and the coefficients $A^{(i)}$ and β_i are

$$\begin{aligned} \beta_1 &= \frac{33 - 2n_f}{12}, \quad \beta_2 = \frac{153 - 19n_f}{24}, \\ A^{(1)} &= \frac{4}{3}, \quad A^{(2)} = \frac{67}{9} - \frac{\pi^2}{3} - \frac{10}{27}n_f + \frac{8}{3}\beta_1 \ln\left(\frac{1}{2}e^{\gamma_E}\right), \end{aligned} \quad (\text{A7})$$

n_f is the number of the quark flavors and γ_E is the Euler constant. We will use the one-loop running coupling constant, i.e. we pick up the four terms in the first line of the expression for the function $s(Q, b)$.

REFERENCES

-
- [1] M. Wirbel, B. Stech, M. Bauer, Z. Phys. C29, 637 (1985); M. Bauer, B. Stech, M. Wirbel, Z. Phys. C34, 103 (1987).

- [2] A. Ali, G. Kramer and C.D. Lü, Phys. Rev. D58, 094009 (1998); Phys. Rev. D59, 014005 (1999); C.D. Lü, Nucl. Phys. Proc. Suppl. 74, 227-230 (1999); Y.-H. Chen, H.-Y. Cheng, B. Tseng, K.-C. Yang, Phys. Rev. D60, 094014 (1999); H.-Y. Cheng and K.-C. Yang, Phys. Rev. D62, 054029 (2000).
- [3] M. Neubert and B. Stech, Adv. Ser. Direct. High Energy Phys. 15, 294-344 (1998) e-Print: hep-ph/9705292; M. Neubert and A.A. Petrov, Phys. Lett. B 519, 50 (2001).
- [4] Y.-Y. Keum, H.-n. Li and A. I. Sanda, Phys. Lett. B504, 6 (2001); Phys. Rev. D63, 054008 (2001); C.-D. Lü, K. Ukai and M.-Z. Yang, Phys. Rev. D63, 074009 (2001); C.-D. Lü and M.-Z. Yang, Eur. Phys. J. C23, 275-287 (2002).
- [5] B.H. Hong and C.D. Lu, Sci. China G49, 357-366 (2006)e-Print: hep-ph/0505020.
- [6] S. Mantry, D. Pirjol, I. W. Stewart, Phys. Rev. D68, 114009 (2003).
- [7] T.Kurimoto, H.-n. Li and A. I. Sanda, Phys. Rev. D67, 054028 (2003); Y.-Y. Keum, T.Kurimoto, H.-n. Li, C.-D. Lu and A. I. Sanda, Phys. Rev. D69, 094018 (2004).
- [8] H. n. Li, Phys. Rev. D **52**, 3958 (1995) [arXiv:hep-ph/9412340]; C. Y. Wu, T. W. Yeh and H. n. Li, Phys. Rev. D **53**, 4982 (1996) [arXiv:hep-ph/9510313].
- [9] C.D. Lu, Eur. Phys. J. C24, 121-126 (2002) e-Print: hep-ph/0112127; Phys. Rev. D68, 097502 (2003). [arXiv:hep-ph/0307040]; Y. Li and C.D. Lu, High Energy Phys. Nucl. Phys. 27, 1062-1066 (2003) e-Print: hep-ph/0305278; J. F. Cheng, D. S. Du and C. D. Lu, Eur. Phys. J. C **45**, 711 (2006) [arXiv:hep-ph/0501082].
- [10] C.-D. Lu, M.-Z. Yang, Eur. Phys. J. C28, 515-523 (2003) e-Print: hep-ph/0212373.
- [11] A. Ali et al., Phys. Rev. D76, 074018 (2007) [arXiv:hep-ph/0703162].
- [12] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D **58**, 114006 (1998) [arXiv:hep-ph/9802409]; T. Feldmann, P. Kroll and B. Stech, Phys. Lett. B **449**, 339 (1999) [arXiv:hep-ph/9812269].
- [13] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33**, 1 (2006).
- [14] P. Ball, G. W. Jones and R. Zwicky, Phys. Rev. D **75**, 054004 (2007) [arXiv:hep-ph/0612081].
- [15] H.-n. Li and B.Melic, Eur.Phys.J.C 11,695(1999).
- [16] A.G.Grozin and M.Neubert, Phys.Rev.D 55,272(1997).
- [17] H.Kawamura, J.Kodaira, C.F.Qiao and K.Tanaka, Phys.Lett.B 523,111(2001);536,344(2002).
- [18] F.Zuo, Z.-H. Li and T. Huang, Phys. Lett. B 641,177(2006) F. Zuo, T. Huang, Chin.Phys.Lett 24, 61-64 (2007).
- [19] S. J. Brodsky, T. Huang and G. P. Lepage, in *Particles and Fields-2*, Proceedings of the Banff Summer Institute, Banff, Alberta, 1981, edited by A. Z. Capri and A. N. Kamal (Plenum, New York, 1983), P143; G. P. Lepage, S. J. Brodsky, T. Huang, and P. B. Mackenize, *ibid.*, p83; T. Huang, in *Proceedings of XXth International Conference on High Energy Physics*, Madison, Wisconsin, 1980, edited by L. Durand and L. G. Pondrom, AIP Conf. Proc. No. 69 (AIP, New York, 1981), p1000.
- [20] CLEO Collaboration, M.Artuso et al., Phys.Rev.Lett. 95,251801(2005);CLEO Collaboration, M.Artuso et al., arXiv: 0704.0629[hep-ex];CLEO Collaboration, T.K.Pedlar et al., arXiv: 0704.0437[hep-ex].
- [21] Y.M. Wang, et al., Eur. Phys. J. C54, 107 (2008), arxiv: 0707.1138.

- [22] <http://hfag.phys.ntu.edu.tw/b2charm/summer2006/index.html>.
- [23] C.D. Lu, K. Ukai, Eur. Phys. J. C28, 305-312 (2003) e-Print: hep-ph/0210206; Y. Li and C.D. Lu, J. Phys. G29, 2115-2124 (2003) e-Print: hep-ph/0304288.
- [24] C.S. Kim, S. Oh and C. Yu, Phys. Lett. B621, 259 (2005); C.-W. Chiang and E.Senaha, Phys. Rev. D 75, 074021(2007).
- [25] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014029 (2005); P. Ball and R. Zwicky, JHEP **0604**, 046 (2006); P. Ball and G. W. Jones, JHEP **0703**, 069 (2007).
- [26] V. L. Chernyak and A.R. Zhitnitsky, Phys. Rept. **112**, 173 (1984); V. M. Braun and I. E. Filyanov, Z. Physik C44, 157 (1989); P. Ball, JHEP **9809**, 005 (1998) [hep-ph/9802394]; V. M. Braun and I. E. Filyanov, Z. Physik C48, 239 (1990); A. R. Zhitnisky, I. R. Zhitnitsky and V. L. Chernyak, Sov. J. Nucl. Phys. **41**, 284(1985), Yad. Fiz. **41**, 445 (1985).
- [27] A.L. Kagan, Phys. Lett. B601, 151-163 (2004); H.-n. Li, S. Mishima, Phys. Rev. D71, 054025 (2005); C.D. Lu, Y.l. Shen, J. Zhu, Eur. Phys. J. C41, 311-317 (2005); Y. Li, C.-D. Lu, Phys. Rev. D73, 014024 (2006); H.W. Huang, etc. Phys. Rev. D73, 014011 (2006)
- [28] V. N. Baier and A. G. Grozin, arXiv:hep-ph/9908365, and references therein.